


Listing of Claims:


1. (Currently Amended) A method of reducing the risk ~~eliminating the onset~~ of Type 1 diabetes in a predisposed human patient by up to 90 percent, comprising the steps of:
identifying a human patient predisposed to Type 1 diabetes ~~patient, wherein Type 1 diabetes is detectable in a patient with autoantibodies to β cell antigens;~~ and
orally administering to the patient an effective amount of a 1α -hydroxy vitamin D compound such that the risk of onset of Type 1 diabetes or Type 1 diabetes symptoms is reduced ~~eliminated~~.
2. (Original) The method of claim 1 wherein the compound is selected from the group consisting of $1\alpha,25$ -dihydroxyvitamin D_3 ($1,25-(OH)_2D_3$), 19-nor- $1,25$ -dihydroxyvitamin D_2 (19-nor- $1,25-(OH)_2D_3$), 24-homo-22-dehydro-22E- $1\alpha,25$ -dihydroxyvitamin D_3 (24-homo-22-dehydro-22E- $1,25-(OH)_2D_3$), $1,25$ -dihydroxy-24(E)-dehydro-24-homo-vitamin D_3 ($1,25-(OH)_2$ -24-homo D_3), 19-nor- $1,25$ -dihydroxy-21-epi-vitamin D_3 (19-nor- $1,25-(OH)_2$ -21-epi- D_3), 1α hydroxy vitamin D_3 or 1α hydroxy vitamin D_2 .
3. (Previously Amended) The method of claim 1 wherein the vitamin D compound is selected from the group consisting of vitamin D compounds with the following formula:

wherein X^1 and X^2 are each selected from the group consisting of hydrogen and acyl; wherein Y^1 and Y^2 are each selected from the group consisting of H, O-aryl, O-alkyl, aryl, and alkyl of 1-4 carbons, taken together to form an alkene having the

structure of B_1



where B_1 and B_2 are selected from the group consisting of H, alkyl of 1-4 carbons and aryl, and can have a β or α configuration; $Z^1=Z^2=H$ or Z^1 and Z^2 together are $=CH_2$; and wherein R is an alkyl, hydroxyalkyl or fluoroalkyl group, or R represents the following side chain:



wherein (a) may have an S or R configuration, R^1 represents hydrogen, hydroxy or O-acyl, R^2 and R^3 are each selected from the group consisting of alkyl, hydroxyalkyl and fluoralkyl, or, when taken together represent the group $-(CH_2)_m$ -wherein m is an integer having a value of from 2 to 5, R^4 is selected from the group consisting of hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl and fluoralkyl, wherein if R^5 is hydroxyl or fluoro, R^4 must be hydrogen or alkyl, R^5 is selected from the group consisting of hydrogen, hydroxy, fluorine, alkyl, hydroxyalkyl and fluoroalkyl, or R^4 and R^5 taken together represent double-bonded oxygen, R^6 and R^7 taken together form a carbon-carbon double bond, R^8 may be H or CH_3 , and

wherein n is an integer having a value of from 1 to 5, and wherein the carbon at any one of positions 20, 22, or 23 in the side chain may be replaced by an O, S, or N atom.

4. (Original) The method of claim 1 wherein the oral administration is via diet.

5. (Original) The method of claim 1 wherein the oral administration is at the concentration of between 0.005 μg to 0.2 μg per kilogram of patient weight per day.

Claims 6 – 10 (Cancelled)


11. (New) A method of reducing the risk of Type 1 diabetes in a predisposed human patient by up to 90 percent, comprising the steps of:

identifying a human patient predisposed to Type 1 diabetes, wherein Type 1 diabetes is detectable in a patient with autoantibodies to glutamic acid decarboxylase; and


orally administering to the patient an effective amount of a 1α -hydroxy vitamin D compound such that the risk of onset of Type 1 diabetes or diabetes symptoms is reduced.

12. (New) The method of claim 11 wherein the compound is selected from the group consisting of $1\alpha,25$ -dihydroxyvitamin D_3 ($1,25\text{-(OH)}_2\text{D}_3$), 19-nor- $1,25$ -dihydroxyvitamin D_2 (19-nor- $1,25\text{-(OH)}_2\text{D}_3$), 24-homo-22-dehydro-22E- $1\alpha,25$ -dihydroxyvitamin D_3 (24-homo-22-dehydro-22E- $1,25\text{-(OH)}_2\text{D}_3$), $1,25$ -dihydroxy-24(E)-dehydro-24-homo-vitamin D_3 ($1,25\text{-(OH)}_2$ -24-homo D_3), 19-nor- $1,25$ -dihydroxy-21-epi-vitamin D_3 (19-nor- $1,25\text{-(OH)}_2$ -21-epi- D_3), 1α hydroxy vitamin D_3 or 1α hydroxy vitamin D_2 .


13. (New) The method of claim 11 wherein the vitamin D compound is selected from the group consisting of vitamin D compounds with the following formula:



wherein X^1 and X^2 are each selected from the group consisting of hydrogen and acyl; wherein Y^1 and Y^2 are each selected from the group consisting of H, 0-aryl, 0-alkyl, aryl, and alkyl of 1-4 carbons, taken together to form an alkene having the structure of B_1



where B_1 and B_2 are selected from the group consisting of H, alkyl of 1-4 carbons and aryl, and can have a β or α configuration; $Z^1=Z^2=H$ or Z^1 and Z^2 together are $=CH_2$; and wherein R is an alkyl, hydroxyalkyl or fluoroalkyl group, or R represents the following side chain:



wherein (a) may have an S or R configuration, R^1 represents hydrogen, hydroxy or O-acyl, R^2 and R^3 are each selected from the group consisting of alkyl, hydroxyalkyl and fluoroalkyl, or, when taken together represent the group $-(CH_2)_m$ -wherein m is an integer having a value of

from 2 to 5, R⁴ is selected from the group consisting of hydrogen, hydroxy, fluorine, O-acyl, alkyl, hydroxyalkyl and fluoroalkyl, wherein if R⁵ is hydroxyl or fluoro, R⁴ must be hydrogen or alkyl, R⁵ is selected from the group consisting of hydrogen, hydroxy, fluorine, alkyl, hydroxyalkyl and fluoroalkyl, or R⁴ and R⁵ taken together represent double-bonded oxygen, R⁶ and R⁷ taken together form a carbon-carbon double bond, R⁸ may be H or CH₃, and wherein n is an integer having a value of from 1 to 5, and wherein the carbon at any one of positions 20, 22, or 23 in the side chain may be replaced by an O, S, or N atom.

14. (New) The method of claim 11 wherein the oral administration is via diet.

15. (New) The method of claim 11 wherein the oral administration is at the concentration of between 0.005 μg to 0.2 μg per kilogram of patient weight per day.